

# $S_3 \times S_3$ acts freely on $S^3 \times S^3$

James F. Davis\*

Department of Mathematics

Indiana University

Bloomington, IN 47405

email: jfdavis@indiana.edu

## Abstract

A free action of the direct product of two copies of the symmetric group on 3 elements on the cartesian product of two copies of the 3-sphere is constructed. This nonlinear action is constructed using surgery. The action provides a counterexample to a conjecture of Lewis made in 1968.

Results of P. A. Smith [10] and J. Milnor [5] show that a dihedral group cannot act freely on a sphere. This implies that if a group  $G$  acts freely on a sphere, then any element of order 2 must be central. Based on this G. Lewis [3] conjectured<sup>1</sup> that if a group  $G$  acts freely on  $S^n \times S^n$ , then any subgroup isomorphic to the Klein 4-group ( $\mathbb{Z}_2 \times \mathbb{Z}_2$ ) must intersect the center non-trivially. (This condition must hold true for a free, linear action.) We will use standard techniques from surgery theory to prove the assertion in the title of this note and thus give a counterexample to Lewis' conjecture.

R. G. Swan [11] constructed a finite  $CW$ -complex  $X$  whose fundamental group is the symmetric group  $S_3$  and whose universal cover  $\tilde{X}$  has the homotopy type of the 3-sphere  $S^3$ . Milnor's result shows that  $X$  does not have the homotopy type of a closed manifold. However, there is the following well-known result, of which we sketch a proof.

---

\*Partially support by an NSF grant.

<sup>1</sup>I thank Alejandro Adem bringing this conjecture to my attention and Peter May for providing background for the proof.

**Proposition.** *Swan's complex  $X$  is a Poincaré complex whose Spivak bundle  $\nu : X \rightarrow BG$  reduces to a stable vector bundle  $\tilde{\nu} : X \rightarrow BO$ .*

*Proof.* Swan's construction (or a spectral sequence argument) shows  $H_3\tilde{X} \rightarrow H_3X$  is isomorphic to  $\mathbb{Z} \xrightarrow{\times 6} \mathbb{Z}$ . Hence the transfer  $H_3X \rightarrow H_3\tilde{X}$  is an isomorphism,  $X$  has a fundamental class, and  $X$  is a Poincaré complex.

A direct analysis of the  $J$ -homomorphism  $\pi_*O \rightarrow \pi_*G$  shows  $\pi_i(G/O) \cong \mathbb{Z}, 0, \mathbb{Z}_2, 0, \mathbb{Z}, 0, \mathbb{Z}_2$  for  $i = 0, 1, 2, 3, 4, 5, 6$  (see [4] for background on surgery classifying spaces). Thus there is a single obstruction in  $H^3(X; \pi_2(G/O)) = \mathbb{Z}_2$  to reduction of the Spivak bundle of  $X$ . The transfer to the 3-fold cover  $Y = \tilde{X}/(12)$  of  $X$  is an injection on mod 2 cohomology, so it suffices to show that  $Y$  has the homotopy type of a closed manifold, namely  $\mathbb{R}P^3$ . Proceed as follows. It is easy to find a map  $f : \mathbb{R}P^3 \rightarrow Y$  which is an isomorphism on  $\pi_1$ . Then  $f$  must have odd degree, since  $f^*(a^3) = f^*(a)^3 \neq 0$ , where  $a \in H^1(Y; \mathbb{Z}_2)$  is the generator. By replacing  $f$  by

$$h : \mathbb{R}P^3 \# S^3 \# \dots \# S^3 \rightarrow Y$$

with the orientation on the  $S^3$  chosen suitably and the map on the  $S^3$ 's given by  $S^3 \xrightarrow{\sim} \tilde{X} \rightarrow Y$ , one obtains the desired homotopy equivalence  $h$ .  $\square$

We need a computational lemma for the proof of the main theorem.

**Lemma.** 1.  $\tilde{K}_0(\mathbb{Z}[S_3 \times S_3]) = 0$ .

2.  $L_2^h(\mathbb{Z}[S_3 \times S_3]) = L_2^p(\mathbb{F}_2[S_3 \times S_3])$ .

*Proof.* By induction theory [1], it suffices to prove that  $\tilde{K}_0$  vanishes for the maximal hyperelementary subgroups:  $\mathbb{Z}_3 \times \mathbb{Z}_3$  and  $S_3 \times \mathbb{Z}_2$ . This result for  $\mathbb{Z}_3 \times \mathbb{Z}_3$  is contained in [8, 8.4]. For  $S_3 \times \mathbb{Z}_2$  we use the cartesian square

$$\begin{array}{ccc} \mathbb{Z}[S_3 \times \mathbb{Z}_2] & \longrightarrow & \mathbb{Z}[S_3] \\ \downarrow & & \downarrow \\ \mathbb{Z}[S_3] & \longrightarrow & \mathbb{F}_2[S_3] \end{array}$$

where the top map and left map send the generator of  $\mathbb{Z}_2$  to  $+1$  and  $-1$  respectively. Thus there is a Mayer-Vietoris exact sequence [6, Chap. 3]

$$K_1(\mathbb{Z}[S_3]) \oplus K_1(\mathbb{Z}[S_3]) \rightarrow K_1(\mathbb{F}_2[S_3]) \xrightarrow{\partial} \tilde{K}_0(\mathbb{Z}[S_3 \times \mathbb{Z}_2]) \rightarrow \tilde{K}_0(\mathbb{Z}[S_3]) \oplus \tilde{K}_0(\mathbb{Z}[S_3])$$

By [11] or [9],  $\tilde{K}_0(\mathbb{Z}[S_3]) = 0$ . Now  $\mathbb{F}_2[S_3] = \mathbb{F}_2[\mathbb{Z}_2] \times M_2(\mathbb{F}_2)$ , so  $K_1(\mathbb{F}_2[S_3])$  has order two, generated by (12). This is in the image of the left hand map, so  $\tilde{K}_0(\mathbb{Z}[S_3 \times \mathbb{Z}_2]) = 0$ . Thus  $\tilde{K}_0(\mathbb{Z}[S_3 \times S_3]) = 0$  and  $L_2^h(\mathbb{Z}[S_3 \times S_3]) = L_2^p(\mathbb{Z}[S_3 \times S_3])$ .

The rational representation theory for  $S_3 \times S_3$  is particularly simple:  $\mathbb{Q}[S_3 \times S_3] = \mathbb{Q}[S_3] \otimes_{\mathbb{Q}} \mathbb{Q}[S_3]$  is a product of matrix rings over the rational numbers. It follows that  $L_2^p(\mathbb{Z}[S_3 \times S_3]) = L_2^p(\hat{\mathbb{Z}}_2[S_3 \times S_3])$  where  $\hat{\mathbb{Z}}_2$  is the 2-adic integers (see [2, Section 4]). Since  $\hat{\mathbb{Z}}_2[S_3 \times S_3]$  is a complete semilocal ring with 2 in its Jacobson radical,  $L_2^p(\hat{\mathbb{Z}}_2[S_3 \times S_3]) = L_2^p(\mathbb{F}_2[S_3 \times S_3])$  (see [13]).

□

**Theorem.**  $S_3 \times S_3$  acts freely and smoothly on  $S^3 \times S^3$ .

*Proof.* By the proposition,  $X$  is a Poincaré complex with a reducible Spivak bundle, so a transversality argument (see [4, Theorem 2.23]) gives a degree one normal map

$$\begin{array}{ccc} \nu_M & \xrightarrow{\hat{f}} & \xi \\ \downarrow & & \downarrow \\ M^3 & \xrightarrow{f} & X. \end{array}$$

There is a surgery obstruction [12, 17D],

$$\sigma_*(f \times f, \hat{f} \times \hat{f}) \in L_6^h(\mathbb{Z}[S_3 \times S_3]) = L_6^p(\mathbb{F}_2[S_3 \times S_3])$$

which vanishes if and only if  $f \times f$  is normally bordant to a homotopy equivalence. By the composition formula and product formula [7]

$$\begin{aligned} \sigma_*(f \times f, \hat{f} \times \hat{f}) &= \sigma_*(\text{Id}_M \times f) + \sigma_*(f \times \text{Id}_M) \\ &= \sigma_*(f, \hat{f}) \otimes (\sigma^*M + \sigma^*X) \in L_6^p(\mathbb{F}_2[S_3 \times S_3]) \end{aligned}$$

with  $\sigma_*(f, \hat{f}) \in L_3^p(\mathbb{F}_2[S_3 \times S_3])$  and the symmetric signatures  $\sigma^*M, \sigma^*X \in L_p^3(\mathbb{F}_2[S_3 \times S_3])$ . But  $L_3^p(\mathbb{F}_2[S_3 \times S_3]) = 0$  since the odd projective  $L$ -theory of a complete semilocal ring vanishes. Thus the integral surgery obstruction is zero and  $X \times X$  has the homotopy type of a closed, smooth manifold  $N$  with fundamental group  $S_3 \times S_3$  and universal cover homotopy equivalent to  $S^3 \times S^3$ .

The surgery exact sequence

$$L_7(\mathbb{Z}) \rightarrow \mathcal{S}(S^3 \times S^3) \rightarrow [S^3 \times S^3, G/O] \rightarrow L_6(\mathbb{Z})$$

shows that  $\mathcal{S}(S^3 \times S^3) = *$ , since  $L_7(\mathbb{Z}) = 0$  and  $[S^3 \times S^3, G/O] \rightarrow L_6(\mathbb{Z})$  is isomorphic to  $\mathbb{Z}_2 \xrightarrow{\text{Id}} \mathbb{Z}_2$ , since there is a framed 6-manifold with non-trivial Kervaire invariant (coincidentally  $S^3 \times S^3$  with the Lie invariant framing). Thus the universal cover of  $N$  is diffeomorphic to  $S^3 \times S^3$  and the deck transformations give the desired action.

□

## References

- [1] Dress, A. W. M., “Induction and structure theorems for orthogonal representations of finite groups,” *Ann. of Math. (2)* **102** (1975), 291–325.
- [2] Hambleton, I. and Taylor, L. R., “A guide to the calculation of surgery obstruction groups,” in *Surgery theory and its applications: surveys presented in honor of C.T.C. Wall*, to appear.
- [3] Lewis, G., “Free actions on  $S^n \times S^n$ ,” *Trans. Amer. Math. Soc.* **132** (1968), 531–540.
- [4] Madsen, I. and Milgram, R. J., *The classifying spaces for surgery and cobordism of manifolds*, *Ann. of Math. Stud.* **92**, Princeton University Press, 1979.
- [5] Milnor, J., “Groups which act on  $S^n$  without fixed points,” *Amer. J. Math.* **79** (1957), 623–630.
- [6] Milnor, John, *Introduction to algebraic K-theory*, *Ann. of Math. Stud.* **72**, Princeton University Press, 1971.
- [7] Ranicki, A. “The algebraic theory of surgery. II. Applications to topology,” *Proc. London Math. Soc. (3)* **40** (1980), 193–283.
- [8] Reiner, I. and Ullom, S., “A Mayer-Vietoris sequence for class groups,” *J. Algebra* **31** (1974), 305–342.

- [9] Reiner, I. and Ullom, S., “Remarks on class groups of integral group rings,” in *Symposia Mathematica, Vol. XIII (Convegno di Gruppi e loro Rappresentazioni, INDAM, Rome, 1972)*, 501–516, Academic Press, 1974.
- [10] Smith, P. A., “Permutable periodic transformations,” *Proc. Nat. Acad. Sci. U. S. A.* **30** (1944), 105–108.
- [11] Swan, R.G., “Periodic resolutions for finite groups,” *Ann. of Math. (2)*, **72** (1960), 267–291.
- [12] Wall, C. T. C., *Surgery on compact manifolds*, Academic Press, 1970.
- [13] Wall, C. T. C., “On the classification of Hermitian forms. III. Complete semilocal rings,” *Invent. Math.* **19** (1973), 59–71.